

At-Home Care Video Transcript

ON-SCREEN TEXT AND MUSIC YOUR GUIDE TO AT-HOME CARE

Learn helpful tips to support your treatment experience with RYBREVANT *FASPRO*[™] or RYBREVANT[®]

EGFR, epidermal growth factor receptor; NSCLC, non–small cell lung cancer. For people with exon 19 deletion or exon 21 L858R substitution mutations.

ON-SCREEN TEXT AND MUSIC THROUGHOUT VIDEO

Logos for RYBREVANT *FASPRO*[™] and RYBREVANT[®] + LAZCLUZE[®]

ON-SCREEN TEXT AND MUSIC

LINDSAY BRAY DOUGHERTY
Certified Registered Nurse Practitioner, Doctor of Nursing Practice

LINDSAY BRAY DOUGHERTY, CRNP, DNP
RYBREVANT *FASPRO*[™] or RYBREVANT[®] is a prescription medicine used to treat adults with non–small cell lung cancer that has spread to other parts of the body or cannot be removed by surgery and has certain abnormal *EGFR* genes. It can be used alone or in combination with other medicines depending on the type of non–small cell lung cancer.

ON-SCREEN TEXT AND MUSIC

Take C.A.R.E. of yourself during treatment

- C Complete oral and/or topical antibiotic treatment, as prescribed
- A Apply a mild, ceramide-based moisturizer (such as La Roche-Posay Lipikar or CeraVe^{*†}) on your face and body (except scalp) at least once a day
- R Remember to wear protective clothing and sunscreen with SPF ≥30 and limit direct exposure to sunlight
- E Ensure you wash both fingernails and toenails with antiseptic cleanser (such as chlorhexidine 4%^{*}) once a day

*These over-the-counter products are available without a prescription.

†All trademarks are property of their respective owners.

LINDSAY BRAY DOUGHERTY, CRNP, DNP
So, you've been prescribed a RYBREVANT *FASPRO*[™]–based or RYBREVANT[®]-based regimen.

I've been treating patients like you for many years now, first as a member of the clinical trial team and now in clinical practice, and I have found that certain tips may help you prepare for your injection or infusion appointments.

Always remember that your healthcare team is there for you throughout your journey to help make your treatment as comfortable as possible. Treatments like RYBREVANT FASPRO™ and RYBREVANT® target certain *EGFR* mutations and can cause certain side effects, including those that affect your skin.

Fortunately, there are things that you can do to help reduce these side effects. Making these steps part of your daily routine may help you and improve your treatment experience.

Some daily, at-home care options may help reduce your risk of side effects such as rash. Things like protecting yourself from the sun and using a mild moisturizer can be incorporated into your daily routine.

ON-SCREEN TEXT AND MUSIC

SUPPORTIVE MEDICATIONS

- Oral antibiotics for skin-related side effects
- If you're taking RYBREVANT FASPRO™ or RYBREVANT® + LAZCLUZE®: medicine to lower risk of blood clots for the first 4 months of treatment

LINDSAY BRAY DOUGHERTY, CRNP, DNP

Your doctor or healthcare team may prescribe you daily pills, such as oral antibiotics, to help reduce the risk of skin-related side effects. For example, I prescribe my patients oral antibiotics for the first 3 months of treatment, based on the COCOON study. If you're taking RYBREVANT FASPRO™ or RYBREVANT® in combination with LAZCLUZE®, you will also be prescribed medicine to prevent blood clots for the first 4 months of your treatment.

Let's take a look at how people can incorporate these steps into their routine. Consult your doctor or healthcare team before performing any at-home care, and talk to them about what is right for you.

PERSON 1

My doctor and I talked about the different pills I'd be taking at the start of treatment, including my antibiotics and medicine to prevent blood clots. We decided that a pill box would help me to stay organized and take my medication every day.

ON-SCREEN TEXT AND MUSIC

Actor portrayal.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

Your healthcare team may prescribe you topical antibiotics to reduce your risk of skin-related side effects. These topical antibiotics may have different application, but let's take a look at one example for your scalp.

ON-SCREEN TEXT AND MUSIC

SCALP CARE

Topical antibiotics for skin-related side effects from Month 4 through Month 12 of treatment

Make sure to tell your doctor or healthcare team about any other medicines you are taking.

ON-SCREEN TEXT AND MUSIC

Actor portrayal.

PERSON 2

After my first 3 months of treatment, my doctor told me it's time to stop taking oral antibiotics and to start applying antibiotic lotion. I like to put it on right before bed every night as part of my nighttime routine. I part my hair down the middle and apply it along the part, and then I do the same a couple inches on either side of the part and above each ear. I put a towel on my pillow to keep the lotion off my sheets.

ON-SCREEN TEXT AND MUSIC

For more details on the recommended timing of oral and/or topical antibiotic regimens, refer to the RYBREVANT *FASPRO*[™] or RYBREVANT[®] Prescribing Information.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

Other skin reactions, like itching and dryness, can occur while taking RYBREVANT *FASPRO*[™]-based or RYBREVANT[®]-based regimens. To help reduce the risk of skin reactions, treat your skin as gently as possible. You can do this by using mild, fragrance-free, and ceramide-based moisturizer, such as La Roche-Posay Lipikar or CeraVe, at least once a day on your face and body.

ON-SCREEN TEXT AND MUSIC

SKIN CARE

- Mild, fragrance-free, ceramide-based moisturizer (such as La Roche-Posay Lipikar or CeraVe^{*†})
- Apply to face and body (except scalp) at least once a day

*All trademarks are property of their respective owners.

†These over-the-counter products are available without a prescription.

Talk to your doctor or healthcare team about what is right for you.

ON-SCREEN TEXT AND MUSIC

Actor portrayal.

PERSON 3

I put on my moisturizer after I get out of the shower in the morning, and again at night before I go to sleep. It's become a nice part of my routine. So, I am going to apply a pump of the lotion on my face, and a couple of pumps to my body. And then I

just blend it in like this. I let it soak in for a minute before putting on my clothes for the day.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

Protecting your skin from the sun is very important while taking RYBREVANT *FASPRO*[™]-based or RYBREVANT[®]-based regimens. It's recommended that you limit sun exposure during and for 2 months after treatment, wear protective clothing, and use sunscreen with an SPF of 30 or greater.

ON-SCREEN TEXT AND MUSIC

SUN PROTECTION

- Limit direct sun exposure
- Wear protective clothing
- Use SPF \geq 30

SPF, sun protection factor.

Talk to your doctor or healthcare team about what is right for you.

ON-SCREEN TEXT AND MUSIC

Actor portrayal.

PERSON 4

I keep my sunscreen by the front door so that I can easily put some on before I walk the dog. I put it on any exposed areas of skin, and as you can see, I'm wearing my hat and sun protection clothing. I check that I'm fully covered, and then I'm good to go!

LINDSAY BRAY DOUGHERTY, CRNP, DNP

In addition, you may also be recommended to use an antiseptic cleanser, which may help to reduce the risk of infection around your nails.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

Your doctor or healthcare team may give you instructions for using a skin cleanser like chlorhexidine 4% on your fingernails and toenails.

ON-SCREEN TEXT AND MUSIC

NAIL CARE

- Antiseptic cleanser (such as chlorhexidine 4%*)
- Wash fingernails and toenails once a day

*This over-the-counter product is available without a prescription.

Talk to your doctor about what's right for you, and refer to the RYBREVANT *FASPRO*[™] or RYBREVANT[®] Prescribing Information.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

I typically tell patients to soak them in the cleanser for 5 minutes. Some of my colleagues suggest soaking a cotton ball in the cleanser and using that to clean the nails. Talk to your doctor or healthcare team to find out what's right for you.

PERSON 2

I've picked up this cleanser that my doctor recommended from the drug store in my neighborhood. I keep it by the sink so that I remember to wash my fingernails and toenails after I get out of the shower.

I make sure to get all the crevices between and around my nails and around my toenails as well. I'll wash my hands with regular soap afterwards, so I don't get the cleanser in my eyes or ears.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

These tips can help your treatment experience with RYBREVANT *FASPRO*[™]-based or RYBREVANT[®]-based regimens by reducing your risk of or severity of side effects.

ON-SCREEN TEXT AND MUSIC

Life is more breathtaking than lung cancer.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

Take care of yourself while on treatment because life is more breathtaking than lung cancer.

LINDSAY BRAY DOUGHERTY, CRNP, DNP

Talk to your doctor or healthcare team about what is right for you.

ON-SCREEN TEXT AND MUSIC

Life is more breathtaking than lung cancer.

ON-SCREEN TEXT AND MUSIC

Take C.A.R.E. of yourself during treatment

- C Complete oral and/or topical antibiotic treatment, as prescribed
- A Apply a mild, ceramide-based moisturizer (such as La Roche-Posay Lipikar or CeraVe^{*†}) on your face and body (except scalp) at least once a day
- R Remember to wear protective clothing and sunscreen with SPF ≥30 and limit direct exposure to sunlight
- E Ensure you wash both fingernails and toenails with antiseptic cleanser (such as chlorhexidine 4%^{*}) once a day

^{*}These over-the-counter products are available without a prescription.

[†]All trademarks are property of their respective owners.

Scan [QR code] for more tips

Data rates may apply.

MUSIC

ON-SCREEN TEXT AND NARRATOR

What are RYBREVANT *FASPRO* (amivantamab and hyaluronidase-lpuj) and RYBREVANT (amivantamab-vmjw)?

RYBREVANT *FASPRO* and RYBREVANT are prescription medicines used to treat adults with non–small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s):

- in combination with LAZCLUZE (lazertinib), a prescription medicine, as a first-line treatment for NSCLC
- in combination with carboplatin and pemetrexed for NSCLC in people whose disease has worsened on or after treatment with an EGFR tyrosine kinase inhibitor (TKI)
- in combination with carboplatin and pemetrexed as a first-line treatment for NSCLC
- alone for the treatment of NSCLC in people whose disease has worsened on or after platinum-based chemotherapy.

Your healthcare provider will perform a test to make sure that RYBREVANT *FASPRO* or RYBREVANT is right for you.

It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE are safe and effective in children.

IMPORTANT SAFETY INFORMATION

Do not take RYBREVANT *FASPRO* if you are allergic to hyaluronidase or any of the ingredients in RYBREVANT *FASPRO*.

Before you receive RYBREVANT *FASPRO*, RYBREVANT or LAZCLUZE, tell your healthcare provider (HCP) about all of your medical conditions, including if you:

- have a history of lung or breathing problems, other than cancer.
- are pregnant or plan to become pregnant. RYBREVANT *FASPRO*, RYBREVANT and LAZCLUZE can harm your unborn baby.

Females who are able to become pregnant:

- Your HCP should do a pregnancy test before you start treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.
- If you are taking RYBREVANT *FASPRO* or RYBREVANT, use effective birth control (contraception) during treatment and for 3 months after your last

dose. If you are taking LAZCLUZE, use effective birth control (contraception) during treatment and for 3 weeks after your last dose.

- Tell your HCP right away if you become pregnant or think you might be pregnant during treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

Males who have female partners who are able to become pregnant:

- You should use effective birth control during treatment and for 3 weeks after your last dose of LAZCLUZE.
- are breastfeeding or plan to breastfeed. It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE pass into your breast milk. Do not breastfeed during treatment and for 3 months after your last dose of RYBREVANT *FASPRO* or RYBREVANT. Do not breastfeed during treatment and for 3 weeks after your last dose of LAZCLUZE.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. LAZCLUZE may affect the way other medicines work, and other medicines may affect how LAZCLUZE works. You should not start or stop any medicine before you talk with your HCP that prescribed LAZCLUZE.

What should I avoid while receiving RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE?

RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause skin reactions. You should limit your time in the sun during and for 2 months after your treatment. Wear protective clothing and use broad-spectrum sunscreen during treatment.

What are the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE?

RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE may cause serious side effects, including:

- **allergic and injection-related reactions.** Allergic and injection-related reactions are common with RYBREVANT *FASPRO* and can be severe or serious. Tell your HCP right away if you get any of the following symptoms during or after your injection: shortness of breath, fever, chills, flushing, chest discomfort, dizziness or lightheadedness, or vomiting.
- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT but can be severe or serious and can include life-threatening (anaphylaxis) allergic reaction. Tell your HCP right away if you get any of the following symptoms during your infusion of RYBREVANT: shortness of breath, difficulty breathing, or wheezing, chest discomfort, swelling of your eyes, lips, or tongue, fever, chills, numbness of your tongue, lips, cheeks, or gums, flushing,

skin rash, hives, or itching, nausea or vomiting, stomach cramps, lightheadedness, dizziness, or fainting, or headache.

- **lung problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause lung problems that can lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your HCP right away if you get any new or worsening lung symptoms, including shortness of breath, cough, or fever.
- **blood clot problems.** RYBREVANT *FASPRO* or RYBREVANT, when given in combination with LAZCLUZE, can cause blood clots in the veins of your legs (deep vein thrombosis) or lungs (pulmonary embolism) that can lead to death. Your HCP may start you on medicine to prevent blood clots for the first 4 months of treatment. Tell your HCP right away if you get any signs and symptoms of blood clots, including swelling, pain or tenderness in the leg, sudden unexplained chest pain, or shortness of breath.
- **skin problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause severe rash, including blisters, peeling, skin pain and sores, redness, raised acne-like bumps, itching, and dry skin. Your HCP may start you on an antibiotic for the first 3 months of treatment followed by an antibiotic lotion for your scalp for the next 9 months. During treatment, you should apply a non-comedogenic (does not clog pores) skin moisturizer (ceramide-based or other types of moisturizers that provide long lasting skin hydration and does not include drying ingredients) on your face and whole body (except scalp) and wash your hands and feet every day with 4% chlorhexidine solution. Tell your HCP right away if you get any skin reactions. Your HCP may treat you with a medicine(s) or send you to see a skin specialist (dermatologist) if you get skin reactions during treatment.
- **liver problems.** LAZCLUZE may cause severe liver problems, including increased liver function tests. Your HCP will do blood tests to check your liver before you start and during treatment. Tell your HCP right away if you get any symptoms of liver problems, including pain on the right side of your stomach area (abdomen), dark or brown (tea colored) urine, or yellowing of your skin or the white part of your eyes.
- **eye problems.** RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause eye problems. Tell your HCP right away if you get symptoms of eye problems including eye pain, inflammation of eyelids, dry eyes, eye redness, blurred vision, changes in vision, itchy eyes, excessive tearing, and sensitivity to light. Your HCP may send you to see an eye specialist (ophthalmologist) if you get new or worsening eye problems during treatment. You should not use contact lenses until your eye symptoms are checked by an HCP.

The most common side effects of RYBREVANT *FASPRO* in combination with LAZCLUZE include: rash, infected skin around the nail, muscle and joint pain, feeling very tired, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, nausea, diarrhea, vomiting, constipation, decreased appetite, headache, and changes in certain blood tests.

The most common side effects seen with intravenous (IV) RYBREVANT, which may be experienced with RYBREVANT *FASPRO*, are listed below:

- **RYBREVANT in combination with LAZCLUZE:** rash, infected skin around the nail, infusion-related reaction, muscle and joint pain, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, unusual feeling in the skin (such as tingling or a crawling feeling), feeling very tired, diarrhea, constipation, COVID-19, bleeding, dry skin, decreased appetite, itchy skin, nausea, and changes in certain blood tests.
- **RYBREVANT in combination with carboplatin and pemetrexed:** rash, infected skin around the nail, infusion-related reaction, feeling very tired, nausea, sores in the mouth, constipation, swelling of hands, ankles, feet, face, or all of your body, decreased appetite, muscle and joint pain, vomiting, COVID-19, and changes in certain blood tests.
- **RYBREVANT given alone:** rash, infusion-related reaction, infected skin around the nail, muscle and joint pain, shortness of breath, nausea, swelling of hands, ankles, feet, face, or all of your body, cough, feeling very tired, sores in the mouth, constipation, vomiting, and changes in certain blood tests.

LAZCLUZE may cause fertility problems in males and females, which may affect your ability to have children. Talk to your HCP if this is a concern for you.

Your HCP may temporarily stop, decrease your dose, or completely stop your treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE if you have serious side effects.

These are not all of the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE:

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LAZCLUZE for a condition for which it was not prescribed. Do not give LAZCLUZE to other people, even if they have the same symptoms that you have. It may harm them.

Please read full Prescribing Information for RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE available on this platform or at this presentation.

ON-SCREEN TEXT

cp-493867v3

Patients taking RYBREVANT *FASPRO* or RYBREVANT can receive personalized 1-on-1 support. Scan the QR code to learn more.

Data rates may apply.

[QR code]

© Johnson & Johnson and its affiliates 2026 05/26 cp-529044v4

[Johnson & Johnson LOGO]