

Debra's Story: The Importance of Shared Decision Making

ON-SCREEN TEXT

The following is a testimonial about a patient receiving RYBREVANT® (amivantamab-vmjw) at the time of filming. RYBREVANT® is indicated in combination with lazertinib for the first-line treatment of adult patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R substitution mutations. Individual results may vary. Please talk with your doctor if you have any questions or to see if RYBREVANT® is right for you.

Important Safety Information

RYBREVANT® is a prescription medicine used to treat adults with non—small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s). RYBREVANT® can cause serious side effects such as infusion-related reactions and lung, blood clot, skin, liver and eye problems. Please see additional Important Safety Information at the end of this video and full Prescribing Information available on this website.

A RYBREVANT® Patient Story

Debra

My name is Debra. I am a manager at a thrift store and I manage 47 volunteers. It is a big part of my life. I enjoy the volunteers. I enjoy the customers. Every time I get in my car and headed there, I'm smiling. I have a place that I am needed and that I enjoy.

ON-SCREEN TEXT AND MUSIC

Debra is a real patient and a volunteer with the SHARE Network, a Johnson & Johnson program made up of people who are dedicated to inspiring others through their personal health journeys and stories of caring. Every patient's experience is unique and Debra's journey is not representative of all patients. Debra has been compensated for her time.

ON-SCREEN TEXT AND MUSIC

Steve

Debra's Husband

Steve

My name is Steve. We met, well you want to talk about that one?

Debra

We met at New Year's party. At a bar. And they all told us not to get married. But we've been married for 42 years.

Steve

42 Years. Yes.

ON-SCREEN TEXT AND MUSIC

Debra's Diagnosis

Debra

I had breathing issues all my life. So if that got worse, I thought it was just because of my age. I thought I had pneumonia. I was coughing a lot. I couldn't stop coughing. I went to my primary doctor and they ordered a chest x-ray. They didn't like what they saw.

They ordered a CT scan, had a PET scan, and that's when they discovered that I had stage 4 cancer in my bones, and I had a mass in my lungs.

I never smoked. My parents smoked, and my husband did smoke, but I never did. I thought when you had cancer of the lungs, you had smoke. It was smoking.

I asked my primary doctor to find out which cancer research place would be best. I didn't want to just pick one out of the hat.

The lung biopsy had to be sent away for genetic makeup, which I ended up having a genetic marker. That would mean a targeted treatment that's better fitted for myself.

ON-SCREEN TEXT AND MUSIC

Debra's Lung Cancer Treatment Journey

Debra

The very first meeting with my doctor, he went through what he saw on all the tests.

ON-SCREEN TEXT AND MUSIC

This is Debra's personal experience.

Steve

Once we got the genetic marker back, he gave us the options, and we decided to go on the route that we're on now.

Debra

...which is RYBREVANT and LAZCLUZE.

Well, he explained, you know, the different side effects that might happen. He kept mentioning itching or a rash, and I have not had that. I have swelling in my legs, and I do have lesions on my head.

I'm on doxycycline. I'm on that every day, twice a day, and I'm on a blood thinner. With my treatments, after the first three weeks, it came much more manageable.

ON-SCREEN TEXT AND MUSIC

Debra's Experience with Treatment

This is Debra's personal experience, individual results may vary.

Debra

After I've gone on the RYBREVANT and LAZCLUZE, I went from an 18% mass in my lung down to 2%. All the lesions and even the ones on my bones had diminished, which is phenomenal. We... when we both heard that, we... contained ourselves inside, but we hugged each other when we left. My husband and I, not my doctor and I. But my husband and I. So... but yeah.

ON-SCREEN TEXT AND MUSIC

RYBREVANT FASPRO™

This refers to how long the injection takes. Actual clinic time may vary. Your care team will also give you additional medications and take time to make sure you're doing well before you leave your visit.

Debra

I'm excited to talk to my doctor and see if RYBREVANT *FASPRO* is an option for me because I've heard that the injection is five-to-seven minutes versus hours for the infusions that I'm on now.

ON-SCREEN TEXT AND MUSIC

Support From Family and Friends

I turned to support from my husband first and foremost, but also from my church. They're keeping me in their prayers. And at the thrift store, I have a ton of volunteers that, you know, are supporting me and also some of my customers.

Steve

One of the things that we decided right away once she did get it is that we were going to do our best to make sure that we keep our life the same. We didn't want to deviate from that at all and not let the cancer or the sickness take over our lives.

ON-SCREEN TEXT AND MUSIC

Debra's Advice to New Patients

Debra

I would say to a new patient that they should just stick with treatment. Get through the preliminary, the first steps, and continue on. Yeah.

Steve

Outlook is really, I feel it's really good. I'm really super happy about it. I don't wake up in the middle of the night worrying about it anymore. It's part of our life and it's just something that we have to work through and deal with and get through, and hopefully everything comes out okay.

Debra

Well, we're planning a trip to Mexico. We don't have the itinerary yet, but for me that's okay. I don't care. I'm just in it for the fun, you know, to—he plans everything out and I just go along.

Steve

It's just to have a long, healthy life and that's to keep doing what we can do as long as we can do it.

Debra

Yep.

ON-SCREEN TEXT AND MUSIC

Logos for RYBREVANT *FASPRO*[™] (amivantamab and hyaluronidase-lpuj)

RYBREVANT *FASPRO*[™] Prescribing Information, Horsham, PA: Janssen Biotech, Inc.
RYBREVANT[®] Prescribing Information, Horsham, PA: Janssen Biotech, Inc.

What are RYBREVANT *FASPRO* (amivantamab and hyaluronidase-lpuj) and RYBREVANT (amivantamab-vmjw)?

RYBREVANT *FASPRO* and RYBREVANT are prescription medicines used to treat adults with non—small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s):

- in combination with LAZCLUZE (lazertinib), a prescription medicine, as a first-line treatment for NSCLC
- in combination with carboplatin and pemetrexed for NSCLC in people whose disease has worsened on or after treatment with an EGFR tyrosine kinase inhibitor (TKI)
- in combination with carboplatin and pemetrexed as a first-line treatment for NSCLC
- alone for the treatment of NSCLC in people whose disease has worsened on or after platinum-based chemotherapy.

Your healthcare provider will perform a test to make sure that RYBREVANT *FASPRO* or RYBREVANT is right for you.

It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE are safe and effective in children.

IMPORTANT SAFETY INFORMATION

Do not take RYBREVANT *FASPRO* if you are allergic to hyaluronidase or any of the ingredients in RYBREVANT *FASPRO*.

Before you receive RYBREVANT FASPRO, RYBREVANT or LAZCLUZE, tell your healthcare provider (HCP) about all of your medical conditions, including if you:

- have a history of lung or breathing problems, other than cancer.
- are pregnant or plan to become pregnant. RYBREVANT FASPRO, RYBREVANT and LAZCLUZE can harm your unborn baby.

Females who are able to become pregnant:

- Your HCP should do a pregnancy test before you start treatment with RYBREVANT FASPRO, RYBREVANT, or LAZCLUZE.
- If you are taking RYBREVANT FASPRO or RYBREVANT, use effective birth control (contraception) during treatment and for 3 months after your last dose. If you are taking LAZCLUZE, use effective birth control (contraception) during treatment and for 3 weeks after your last dose.
- Tell your HCP right away if you become pregnant or think you might be pregnant during treatment with RYBREVANT FASPRO, RYBREVANT, or LAZCLUZE.

Males who have female partners who are able to become pregnant:

- You should use effective birth control during treatment and for 3 weeks after your last dose of LAZCLUZE.
- are breastfeeding or plan to breastfeed. It is not known if RYBREVANT FASPRO, RYBREVANT, or LAZCLUZE pass into your breast milk. Do not breastfeed during treatment and for 3 months after your last dose of RYBREVANT FASPRO or RYBREVANT. Do not breastfeed during treatment and for 3 weeks after your last dose of LAZCLUZE.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. LAZCLUZE may affect the way other medicines work, and other medicines may affect how LAZCLUZE works. You should not start or stop any medicine before you talk with your HCP that prescribed LAZCLUZE.

What should I avoid while receiving RYBREVANT FASPRO, RYBREVANT, and/or LAZCLUZE?

RYBREVANT FASPRO, RYBREVANT, and LAZCLUZE can cause skin reactions. You should limit your time in the sun during and for 2 months after your treatment. Wear protective clothing and use broad-spectrum sunscreen during treatment.

What are the possible side effects of RYBREVANT FASPRO, RYBREVANT, or LAZCLUZE?

RYBREVANT FASPRO, RYBREVANT, and/or LAZCLUZE may cause serious side effects, including:

- **allergic and injection-related reactions.** Allergic and injection-related reactions are common with RYBREVANT FASPRO and can be severe or serious. Tell your HCP right away if you get any of the following symptoms during or after your injection: shortness of breath, fever, chills, flushing, chest discomfort, dizziness or lightheadedness, or vomiting.

- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT but can be severe or serious and can include life-threatening (anaphylaxis) allergic reaction. Tell your HCP right away if you get any of the following symptoms during your infusion of RYBREVANT: shortness of breath, difficulty breathing, or wheezing, chest discomfort, swelling of your eyes, lips, or tongue, fever, chills, numbness of your tongue, lips, cheeks, or gums, flushing, skin rash, hives, or itching, nausea or vomiting, stomach cramps, lightheadedness, dizziness, or fainting, or headache.
- **lung problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause lung problems that can lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your HCP right away if you get any new or worsening lung symptoms, including shortness of breath, cough, or fever.
- **blood clot problems.** RYBREVANT *FASPRO* or RYBREVANT, when given in combination with LAZCLUZE, can cause blood clots in the veins of your legs (deep vein thrombosis) or lungs (pulmonary embolism) that can lead to death. Your HCP may start you on medicine to prevent blood clots for the first 4 months of treatment. Tell your HCP right away if you get any signs and symptoms of blood clots, including swelling, pain or tenderness in the leg, sudden unexplained chest pain, or shortness of breath.
- **skin problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause severe rash, including blisters, peeling, skin pain and sores, redness, raised acne-like bumps, itching, and dry skin. Your HCP may start you on an antibiotic for the first 3 months of treatment followed by an antibiotic lotion for your scalp for the next 9 months. During treatment, you should apply a non-comedogenic (does not clog pores) skin moisturizer (ceramide-based or other types of moisturizers that provide long lasting skin hydration and does not include drying ingredients) on your face and whole body (except scalp) and wash your hands and feet every day with 4% chlorhexidine solution. Tell your HCP right away if you get any skin reactions. Your HCP may treat you with a medicine(s) or send you to see a skin specialist (dermatologist) if you get skin reactions during treatment.
- **liver problems.** LAZCLUZE may cause severe liver problems, including increased liver function tests. Your HCP will do blood tests to check your liver before you start and during treatment. Tell your HCP right away if you get any symptoms of liver problems, including pain on the right side of your stomach area (abdomen), dark or brown (tea colored) urine, or yellowing of your skin or the white part of your eyes.
- **eye problems.** RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause eye problems. Tell your HCP right away if you get symptoms of eye problems including eye pain, inflammation of eyelids, dry eyes, eye redness, blurred vision, changes in vision, itchy eyes, excessive tearing, and sensitivity to light. Your HCP may send you to see an eye specialist (ophthalmologist) if you get new or worsening eye problems during treatment. You should not use contact lenses until your eye symptoms are checked by an HCP.

The most common side effects of RYBREVANT *FASPRO* in combination with LAZCLUZE include: rash, infected skin around the nail, muscle and joint pain, feeling very tired, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body,

nausea, diarrhea, vomiting, constipation, decreased appetite, headache, and changes in certain blood tests.

The most common side effects seen with intravenous (IV) RYBREVANT, which may be experienced with RYBREVANT FASPRO, are listed below:

- **RYBREVANT in combination with LAZCLUZE:** rash, infected skin around the nail, infusion-related reaction, muscle and joint pain, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, unusual feeling in the skin (such as tingling or a crawling feeling), feeling very tired, diarrhea, constipation, COVID-19, bleeding, dry skin, decreased appetite, itchy skin, nausea, and changes in certain blood tests.
- **RYBREVANT in combination with carboplatin and pemetrexed:** rash, infected skin around the nail, infusion-related reaction, feeling very tired, nausea, sores in the mouth, constipation, swelling of hands, ankles, feet, face, or all of your body, decreased appetite, muscle and joint pain, vomiting, COVID-19, and changes in certain blood tests.
- **RYBREVANT given alone:** rash, infusion-related reaction, infected skin around the nail, muscle and joint pain, shortness of breath, nausea, swelling of hands, ankles, feet, face, or all of your body, cough, feeling very tired, sores in the mouth, constipation, vomiting, and changes in certain blood tests.

LAZCLUZE may cause fertility problems in males and females, which may affect your ability to have children. Talk to your HCP if this is a concern for you.

Your HCP may temporarily stop, decrease your dose, or completely stop your treatment with RYBREVANT FASPRO, RYBREVANT, or LAZCLUZE if you have serious side effects.

These are not all of the possible side effects of RYBREVANT FASPRO, RYBREVANT, or LAZCLUZE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of RYBREVANT FASPRO, RYBREVANT, or LAZCLUZE:

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LAZCLUZE for a condition for which it was not prescribed. Do not give LAZCLUZE to other people, even if they have the same symptoms that you have. It may harm them.

Please read full Prescribing Information for [RYBREVANT FASPRO](#), [RYBREVANT](#), and [LAZCLUZE](#) available on this platform or at this presentation.

cp-493867v3