

Heather's Story: Finding Resilience Through the Journey

ON-SCREEN TEXT

The following is a testimonial about a patient receiving RYBREVANT® (amivantamab-vmjw) at the time of filming. RYBREVANT® is indicated in combination with carboplatin and pemetrexed as a first-line treatment for people who have locally advanced or metastatic non–small cell lung cancer with *EGFR* exon 20 insertion mutations. Individual results may vary. Please talk with your doctor if you have any questions or to see if RYBREVANT® is right for you.

Important Safety Information

RYBREVANT® is a prescription medicine used to treat adults with non–small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (*EGFR*) gene(s). RYBREVANT® can cause serious side effects such as infusion-related reactions and lung, blood clot, skin and eye problems. Please see additional Important Safety Information at the end of this video and full Prescribing Information available on this website.

ON-SCREEN TEXT AND MUSIC

Heather

My name is Heather, and I'm a mother of two children, age five and eight, two boys, and I've been married for ten years.

I'm a working mom. I work in the music industry making merchandise for artists, one of the biggest record labels in the world.

A RYBREVANT® Patient Story

Heather

Heather is a real patient and a volunteer with the SHARE Network, a Johnson & Johnson program made up of people who are dedicated to inspiring others through their personal health journeys and stories of caring. Every patient's experience is unique, and Heather's journey is not representative of all patients. Heather has been compensated for her time.

ON-SCREEN TEXT AND MUSIC

My Diagnosis

Heather

I had a cough that wouldn't go away. And as a mother of young children, a cough is something that I've almost always had. It persisted for months and months. And then eventually, I realized that it seemed a little bit more serious, almost pneumonia like.

So I ended up going to get checked out for pneumonia. The results came in, and it said that I had metastatic cancer. And the cancer had spread from my lungs to my

bones, to my liver, to my lymph nodes, and at that point, I could not believe that not only did I have cancer, but I had it everywhere.

ON-SCREEN TEXT AND MUSIC

Heather received genetic testing prior to this appointment.

Heather

The day I got to the hospital, I was there for about eight hours that day, and they went from hi, nice to meet you, to we figured out your genetic mutation, we have a plan of medication that was recently FDA-approved just about two weeks before I got there, and they said, we want to be aggressive, we want to be swift. Are you with us? And I said absolutely yes.

My oncologist said I want you to be the new statistic. We're going to change these numbers. I trusted my doctor, and I said, I will do, I'll drive as hard as you want me to drive.

ON-SCREEN TEXT AND MUSIC

The Lung Cancer

Treatment Journey

Heather

I was thinking when I started out becoming a cancer patient, that the way the road would go would be that I would do a treatment regimen for a certain amount of time, and then I would go into a remission or there would be like a clean ending, and I realized that living with metastatic cancer and a cancer that is treatable, not curable, the road looks a little different, and I have to be willing to go with where the road takes me.

When I got my diagnosis that I had lung cancer, people said, I had to explain that I'm not a smoker, I've never smoked in my life. They say were you around anyone that smoked, and I said no, it doesn't work like that. It's a genetic mutation. Well, what caused a genetic mutation? Nobody knows.

I think that lung cancer has a perception of being difficult to treat, that it's some, you know, some sort of impossible situation. That's just not the truth anymore. I feel like there's a lot of advancements happening, specifically in lung cancer, that they're making some great breakthroughs with medications and targeted therapies like RYBREVANT®.

And the biggest takeaway I want people to understand is that even if you have stage 4, you have options to live a very full life if you manage it properly. You know, by taking medication for it and managing any side effects those might cause, it's so worth it.

ON-SCREEN TEXT AND MUSIC

My Experience with Treatment

When I started on my treatment plan, I was going to be doing three medications, pemetrexed, RYBREVANT[®], and carboplatin.

It was interesting to me when I got my treatment plan for RYBREVANT[®] to understand what it was that I was taking, that it was a smart drug that would follow the cancer from where it started and only go to the places that it needed to where it had spread.

ON-SCREEN TEXT AND MUSIC

This is Heather's personal experience.

Heather

They would scan me after, you know, the first three treatments, and my cancer had shrunk over 50% in everywhere that it had been. And I could not believe that a medicine really would work that effectively. But not only working on my lung, but it worked on all the other parts of my body.

And I continued on, and the cancer is totally inactive now that I completed my treatment. So it worked. I cannot believe that it's so effective. People need to be educated about, you know, the new technology and how smart these drugs are becoming. It can really save your life.

Once I'm done with treatment, I go home and I pick my kids up from school. And it's pretty amazing to me that I can go and get medicine and still be in shape to pick up my kids.

I've heard that there is a RYBREVANT[®] injection, would take five-to-seven minutes.

ON-SCREEN TEXT AND MUSIC

This refers to how long the injection takes. Actual clinic time may vary. Your care team will also give you additional medications and take time to make sure you're doing well before you leave your visit.

Heather

That would be so incredible for me. Because right now, I'm spending the whole day at the hospital, and it's very hard for me to pick up my kids and be there perfectly on time. And to get those hours back with an injection of my medicine would be such a game changer for me.

It's my goal to live to be an old lady and to be there for my children and my husband. And it may not be a straight road. It might take some twists and turns, but the point is that you have to stay on the road.

ON-SCREEN TEXT AND MUSIC

Logos for RYBREVANT FASPRO[™] (amivantamab and hyaluronidase-lpluj)
RYBREVANT FASPRO[™] Prescribing Information, Horsham, PA Janssen Biotech, Inc.

ON-SCREEN TEXT AND NARRATOR

What are RYBREVANT *FASPRO* (amivantamab and hyaluronidase-lpuj) and RYBREVANT (amivantamab-vmjw)?

RYBREVANT *FASPRO* and RYBREVANT are prescription medicines used to treat adults with non–small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s):

- in combination with LAZCLUZE (lazertinib), a prescription medicine, as a first-line treatment for NSCLC
- in combination with carboplatin and pemetrexed for NSCLC in people whose disease has worsened on or after treatment with an EGFR tyrosine kinase inhibitor (TKI)
- in combination with carboplatin and pemetrexed as a first-line treatment for NSCLC
- alone for the treatment of NSCLC in people whose disease has worsened on or after platinum-based chemotherapy.

Your healthcare provider will perform a test to make sure that RYBREVANT *FASPRO* or RYBREVANT is right for you.

It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE are safe and effective in children.

IMPORTANT SAFETY INFORMATION

Do not take RYBREVANT *FASPRO* if you are allergic to hyaluronidase or any of the ingredients in RYBREVANT *FASPRO*.

Before you receive RYBREVANT *FASPRO*, RYBREVANT or LAZCLUZE, tell your healthcare provider (HCP) about all of your medical conditions, including if you:

- have a history of lung or breathing problems, other than cancer.
- are pregnant or plan to become pregnant. RYBREVANT *FASPRO*, RYBREVANT and LAZCLUZE can harm your unborn baby.

Females who are able to become pregnant:

- Your HCP should do a pregnancy test before you start treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.
- If you are taking RYBREVANT *FASPRO* or RYBREVANT, use effective birth control (contraception) during treatment and for 3 months after your last dose. If you are taking LAZCLUZE, use effective birth control (contraception) during treatment and for 3 weeks after your last dose.
- Tell your HCP right away if you become pregnant or think you might be pregnant during treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

Males who have female partners who are able to become pregnant:

- You should use effective birth control during treatment and for 3 weeks after your last dose of LAZCLUZE.

- are breastfeeding or plan to breastfeed. It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE pass into your breast milk. Do not breastfeed during treatment and for 3 months after your last dose of RYBREVANT *FASPRO* or RYBREVANT. Do not breastfeed during treatment and for 3 weeks after your last dose of LAZCLUZE.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. LAZCLUZE may affect the way other medicines work, and other medicines may affect how LAZCLUZE works. You should not start or stop any medicine before you talk with your HCP that prescribed LAZCLUZE.

What should I avoid while receiving RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE?

RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause skin reactions. You should limit your time in the sun during and for 2 months after your treatment. Wear protective clothing and use broad-spectrum sunscreen during treatment.

What are the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE?

RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE may cause serious side effects, including:

- **allergic and injection-related reactions.** Allergic and injection-related reactions are common with RYBREVANT *FASPRO* and can be severe or serious. Tell your HCP right away if you get any of the following symptoms during or after your injection: shortness of breath, fever, chills, flushing, chest discomfort, dizziness or lightheadedness, or vomiting.
- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT but can be severe or serious and can include life-threatening (anaphylaxis) allergic reaction. Tell your HCP right away if you get any of the following symptoms during your infusion of RYBREVANT: shortness of breath, difficulty breathing, or wheezing, chest discomfort, swelling of your eyes, lips, or tongue, fever, chills, numbness of your tongue, lips, cheeks, or gums, flushing, skin rash, hives, or itching, nausea or vomiting, stomach cramps, lightheadedness, dizziness, or fainting, or headache.
- **lung problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause lung problems that can lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your HCP right away if you get any new or worsening lung symptoms, including shortness of breath, cough, or fever.
- **blood clot problems.** RYBREVANT *FASPRO* or RYBREVANT, when given in combination with LAZCLUZE, can cause blood clots in the veins of your legs (deep vein thrombosis) or lungs (pulmonary embolism) that can lead to death. Your HCP may start you on medicine to prevent blood clots for the first 4 months of treatment. Tell your HCP right away if you get any signs and symptoms of blood clots, including swelling, pain or tenderness in the leg, sudden unexplained chest pain, or shortness of breath.
- **skin problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause

severe rash, including blisters, peeling, skin pain and sores, redness, raised acne-like bumps, itching, and dry skin. Your HCP may start you on an antibiotic for the first 3 months of treatment followed by an antibiotic lotion for your scalp for the next 9 months. During treatment, you should apply a non-comedogenic (does not clog pores) skin moisturizer (ceramide-based or other types of moisturizers that provide long lasting skin hydration and does not include drying ingredients) on your face and whole body (except scalp) and wash your hands and feet every day with 4% chlorhexidine solution. Tell your HCP right away if you get any skin reactions. Your HCP may treat you with a medicine(s) or send you to see a skin specialist (dermatologist) if you get skin reactions during treatment.

- **liver problems.** LAZCLUZE may cause severe liver problems, including increased liver function tests. Your HCP will do blood tests to check your liver before you start and during treatment. Tell your HCP right away if you get any symptoms of liver problems, including pain on the right side of your stomach area (abdomen), dark or brown (tea colored) urine, or yellowing of your skin or the white part of your eyes.
- **eye problems.** RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause eye problems. Tell your HCP right away if you get symptoms of eye problems including eye pain, inflammation of eyelids, dry eyes, eye redness, blurred vision, changes in vision, itchy eyes, excessive tearing, and sensitivity to light. Your HCP may send you to see an eye specialist (ophthalmologist) if you get new or worsening eye problems during treatment. You should not use contact lenses until your eye symptoms are checked by an HCP.

The most common side effects of RYBREVANT *FASPRO* in combination with LAZCLUZE include: rash, infected skin around the nail, muscle and joint pain, feeling very tired, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, nausea, diarrhea, vomiting, constipation, decreased appetite, headache, and changes in certain blood tests.

The most common side effects seen with intravenous (IV) RYBREVANT, which may be experienced with RYBREVANT *FASPRO*, are listed below:

- **RYBREVANT in combination with LAZCLUZE:** rash, infected skin around the nail, infusion-related reaction, muscle and joint pain, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, unusual feeling in the skin (such as tingling or a crawling feeling), feeling very tired, diarrhea, constipation, COVID-19, bleeding, dry skin, decreased appetite, itchy skin, nausea, and changes in certain blood tests.
- **RYBREVANT in combination with carboplatin and pemetrexed:** rash, infected skin around the nail, infusion-related reaction, feeling very tired, nausea, sores in the mouth, constipation, swelling of hands, ankles, feet, face, or all of your body, decreased appetite, muscle and joint pain, vomiting, COVID-19, and changes in certain blood tests.
- **RYBREVANT given alone:** rash, infusion-related reaction, infected skin around the nail, muscle and joint pain, shortness of breath, nausea, swelling of hands, ankles, feet, face, or all of your body, cough, feeling very tired, sores in the mouth, constipation, vomiting, and changes in certain blood tests.

LAZCLUZE may cause fertility problems in males and females, which may affect your ability to have children. Talk to your HCP if this is a concern for you.

Your HCP may temporarily stop, decrease your dose, or completely stop your treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE if you have serious side effects.

These are not all of the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE:

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LAZCLUZE for a condition for which it was not prescribed. Do not give LAZCLUZE to other people, even if they have the same symptoms that you have. It may harm them.

Please read full Prescribing Information for [RYBREVANT *FASPRO*](#), [RYBREVANT](#), and [LAZCLUZE](#) available on this platform or at this presentation.

cp-493867v3