

# Jeanne's Story: Leaning on Your Support System

## ON-SCREEN TEXT

The following is a testimonial about a patient receiving RYBREVANT® (amivantamab-vmjw) at the time of filming. RYBREVANT® is a prescription medicine used to treat adults with non–small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s). Individual results may vary. Please talk with your doctor if you have any questions or to see if RYBREVANT® is right for you.

## Important Safety Information

RYBREVANT® can cause serious side effects such as infusion-related reactions and lung, blood clot, skin, liver and eye problems. Please see additional Important Safety Information at the end of this video and full Prescribing Information available on this website.

## ON-SCREEN TEXT AND MUSIC

### A RYBREVANT® Patient Story

#### Jeanne

My name is Jeanne and I come from Minnesota. I grew up in southern Minnesota, and moved to the Twin Cities to go to college. Met my husband there and have two wonderful children.

I worked in public relations for several years. Then I volunteered at school and other organizations until I was hired by one of the programs, and I retired in 2024.

## ON-SCREEN TEXT AND MUSIC

Jeanne is a real patient and a volunteer with the SHARE Network, a Johnson & Johnson program made up of people who are dedicated to inspiring others through their personal health journeys and stories of caring. Every patient's experience is unique, and Jeanne's journey is not representative of all patients. Jeanne has been compensated for her time.

#### Jeanne

Family is very important. We have our family and we have what we call our chosen family, which are friends that we have known since college.

I love traveling internationally. So I go to those places mostly with my daughter. She's my travel buddy.

## ON-SCREEN TEXT AND MUSIC

### Symptoms & Diagnosis

#### Jeanne

So I had virtually no symptoms of my disease prior to getting really sick. My husband and I, we set off on a 33-day cruise. We had a wonderful time, and I really had no symptoms. We got home, and two days later I got up, took my shower, and I was exhausted and I went back to bed. And that's where I was through December.

I did see a doctor who treated me for pneumonia. I didn't feel better so she treated me with a more heavy-duty antibiotic and still not, not getting better. So, I had a call in to my primary doctor. They said, you know, head to the emergency room, and they sent me down for a CT, and then I knew something was up.

Finally, these two doctors came into the room in the Emergency Department and they said it looks like you have cancer in your lungs and all over in your body. He said we will keep you as comfortable as possible.

I had lesions on many bones, on my skull, on my spine, all over in my right lung. There was a huge tumor, huge neoplasm on the bronchus. And I didn't ever think that somebody that didn't smoke could get lung cancer. I've since found out a lot differently. Like my doctor said, if you have lungs, you can get cancer.

The saving grace is that one of the first things that my doctor ordered from the emergency room was the genetic testing. He said that I had what was called an EGFR mutation and that there was a drug that I could take for it.

## ON-SCREEN TEXT AND MUSIC

### The Lung Cancer

### Treatment Journey

#### Jeanne

And he suggested that I try this new combination called RYBREVANT and LAZCLUZE because it had shown great results and he thought it would work for me.

## ON-SCREEN TEXT AND MUSIC

This is Jeanne's personal experience.

#### Jeanne

It's a new drug, but they've worked very hard to prepare the patient to handle the possible reactions and side effects over the course of the clinical studies. He said you will take doxycycline twice a day. I started out by using the antibacterial lotion on my fingernails and my toenails. I have to use sunscreen. You'll use this moisturizer every day. I was on a blood thinner.

I believe that symptoms are manageable, especially if you're prepared from the very beginning and you follow those protocols. After only three months on the drug, the cancer was gone. Even the metastases on my skull and my arms and hip were gone. We all cried. We were so amazed. We were expecting good news, and so we were just ecstatic.

I've been on the RYBREVANT/LAZCLUZE treatment getting biweekly infusion of RYBREVANT.

## ON-SCREEN TEXT AND MUSIC

### RYBREVANT *FASPRO*<sup>™</sup>

This refers to how long the injection takes. Actual clinic time may vary. Your care team will also give you additional medications and take time to make sure you're doing well before you leave your visit.

#### Jeanne

My doctor told me that RYBREVANT *FASPRO* injection, a subcutaneous form, which takes five-to-seven minutes, which will reduce the time that I'm in the oncology infusion room by hours. I told him, put me at the top of the list, I am ready to go.

## ON-SCREEN TEXT AND MUSIC

### My Takeaway

#### Jeanne

I wanted to share my story because I can help somebody else learn I'm not alone and that there are things I can do to help myself get the care that I need.

I learned really early on that it's important to advocate for your health, especially with lung cancer striking so many non-smokers. A simple cough is all I had as a symptom and it was dismissed as possibly allergies.

My support network is varied. My husband, my family, my friends and my oncology team all working together. I thank them for their support and I hope to live long enough to take care of them when they need help.

I know that I'll never hear "cured" because you don't cure stage 4 cancer, but if I can buy some time here, whether it's three years, five years, ten years, I'll take it. I'm looking forward to be able to do everything that I used to do as I'm kind of slowly working my way back. That means a lot to be able to look ahead.

## ON-SCREEN TEXT

Logos for RYBREVANT *FASPRO*<sup>™</sup> (amivantamab and hyaluronidase-lpuj)

RYBREVANT *FASPRO*<sup>™</sup> Prescribing Information, Horsham, PA: Janssen Biotech, Inc.  
RYBREVANT<sup>®</sup> Prescribing Information, Horsham, PA: Janssen Biotech, Inc.

## ON-SCREEN TEXT AND NARRATOR

### **What are RYBREVANT *FASPRO* (amivantamab and hyaluronidase-lpuj) and RYBREVANT (amivantamab-vmjw)?**

RYBREVANT *FASPRO* and RYBREVANT are prescription medicines used to treat adults with non-small cell lung cancer (NSCLC) that has spread to other parts of the

body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s):

- in combination with LAZCLUZE (lazertinib), a prescription medicine, as a first-line treatment for NSCLC
- in combination with carboplatin and pemetrexed for NSCLC in people whose disease has worsened on or after treatment with an EGFR tyrosine kinase inhibitor (TKI)
- in combination with carboplatin and pemetrexed as a first-line treatment for NSCLC
- alone for the treatment of NSCLC in people whose disease has worsened on or after platinum-based chemotherapy.

Your healthcare provider will perform a test to make sure that RYBREVANT *FASPRO* or RYBREVANT is right for you.

It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE are safe and effective in children.

### **IMPORTANT SAFETY INFORMATION**

**Do not take RYBREVANT *FASPRO* if you are allergic to hyaluronidase or any of the ingredients in RYBREVANT *FASPRO*.**

**Before you receive RYBREVANT *FASPRO*, RYBREVANT or LAZCLUZE, tell your healthcare provider (HCP) about all of your medical conditions, including if you:**

- have a history of lung or breathing problems, other than cancer.
- are pregnant or plan to become pregnant. RYBREVANT *FASPRO*, RYBREVANT and LAZCLUZE can harm your unborn baby.

#### **Females who are able to become pregnant:**

- Your HCP should do a pregnancy test before you start treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.
- If you are taking RYBREVANT *FASPRO* or RYBREVANT, use effective birth control (contraception) during treatment and for 3 months after your last dose. If you are taking LAZCLUZE, use effective birth control (contraception) during treatment and for 3 weeks after your last dose.
- Tell your HCP right away if you become pregnant or think you might be pregnant during treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

#### **Males who have female partners who are able to become pregnant:**

- You should use effective birth control during treatment and for 3 weeks after your last dose of LAZCLUZE.
- are breastfeeding or plan to breastfeed. It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE pass into your breast milk. Do not breastfeed during treatment and for 3 months after your last dose of RYBREVANT *FASPRO* or RYBREVANT. Do not breastfeed during treatment and for 3 weeks after your last dose of LAZCLUZE.

**Tell your HCP about all the medicines you take**, including prescription and over-the-

counter medicines, vitamins, and herbal supplements. LAZCLUZE may affect the way other medicines work, and other medicines may affect how LAZCLUZE works. You should not start or stop any medicine before you talk with your HCP that prescribed LAZCLUZE.

**What should I avoid while receiving RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE?**

RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause skin reactions. You should limit your time in the sun during and for 2 months after your treatment. Wear protective clothing and use broad-spectrum sunscreen during treatment.

**What are the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE?**

**RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE may cause serious side effects, including:**

- **allergic and injection-related reactions.** Allergic and injection-related reactions are common with RYBREVANT *FASPRO* and can be severe or serious. Tell your HCP right away if you get any of the following symptoms during or after your injection: shortness of breath, fever, chills, flushing, chest discomfort, dizziness or lightheadedness, or vomiting.
- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT but can be severe or serious and can include life-threatening (anaphylaxis) allergic reaction. Tell your HCP right away if you get any of the following symptoms during your infusion of RYBREVANT: shortness of breath, difficulty breathing, or wheezing, chest discomfort, swelling of your eyes, lips, or tongue, fever, chills, numbness of your tongue, lips, cheeks, or gums, flushing, skin rash, hives, or itching, nausea or vomiting, stomach cramps, lightheadedness, dizziness, or fainting, or headache.
- **lung problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause lung problems that can lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your HCP right away if you get any new or worsening lung symptoms, including shortness of breath, cough, or fever.
- **blood clot problems.** RYBREVANT *FASPRO* or RYBREVANT, when given in combination with LAZCLUZE, can cause blood clots in the veins of your legs (deep vein thrombosis) or lungs (pulmonary embolism) that can lead to death. Your HCP may start you on medicine to prevent blood clots for the first 4 months of treatment. Tell your HCP right away if you get any signs and symptoms of blood clots, including swelling, pain or tenderness in the leg, sudden unexplained chest pain, or shortness of breath.
- **skin problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause severe rash, including blisters, peeling, skin pain and sores, redness, raised acne-like bumps, itching, and dry skin. Your HCP may start you on an antibiotic for the first 3 months of treatment followed by an antibiotic lotion for your scalp for the next 9 months. During treatment, you should apply a non-comedogenic (does not clog pores) skin moisturizer (ceramide-based or other types of moisturizers that provide long lasting skin hydration and does not include drying ingredients) on

your face and whole body (except scalp) and wash your hands and feet every day with 4% chlorhexidine solution. Tell your HCP right away if you get any skin reactions. Your HCP may treat you with a medicine(s) or send you to see a skin specialist (dermatologist) if you get skin reactions during treatment.

- **liver problems.** LAZCLUZE may cause severe liver problems, including increased liver function tests. Your HCP will do blood tests to check your liver before you start and during treatment. Tell your HCP right away if you get any symptoms of liver problems, including pain on the right side of your stomach area (abdomen), dark or brown (tea colored) urine, or yellowing of your skin or the white part of your eyes.
- **eye problems.** RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause eye problems. Tell your HCP right away if you get symptoms of eye problems including eye pain, inflammation of eyelids, dry eyes, eye redness, blurred vision, changes in vision, itchy eyes, excessive tearing, and sensitivity to light. Your HCP may send you to see an eye specialist (ophthalmologist) if you get new or worsening eye problems during treatment. You should not use contact lenses until your eye symptoms are checked by an HCP.

**The most common side effects of RYBREVANT *FASPRO* in combination with LAZCLUZE include:** rash, infected skin around the nail, muscle and joint pain, feeling very tired, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, nausea, diarrhea, vomiting, constipation, decreased appetite, headache, and changes in certain blood tests.

**The most common side effects seen with intravenous (IV) RYBREVANT, which may be experienced with RYBREVANT *FASPRO*, are listed below:**

- **RYBREVANT in combination with LAZCLUZE:** rash, infected skin around the nail, infusion-related reaction, muscle and joint pain, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, unusual feeling in the skin (such as tingling or a crawling feeling), feeling very tired, diarrhea, constipation, COVID-19, bleeding, dry skin, decreased appetite, itchy skin, nausea, and changes in certain blood tests.
- **RYBREVANT in combination with carboplatin and pemetrexed:** rash, infected skin around the nail, infusion-related reaction, feeling very tired, nausea, sores in the mouth, constipation, swelling of hands, ankles, feet, face, or all of your body, decreased appetite, muscle and joint pain, vomiting, COVID-19, and changes in certain blood tests.
- **RYBREVANT given alone:** rash, infusion-related reaction, infected skin around the nail, muscle and joint pain, shortness of breath, nausea, swelling of hands, ankles, feet, face, or all of your body, cough, feeling very tired, sores in the mouth, constipation, vomiting, and changes in certain blood tests.

LAZCLUZE may cause fertility problems in males and females, which may affect your ability to have children. Talk to your HCP if this is a concern for you.

Your HCP may temporarily stop, decrease your dose, or completely stop your treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE if you have serious side effects.

These are not all of the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**General information about the safe and effective use of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE:**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LAZCLUZE for a condition for which it was not prescribed. Do not give LAZCLUZE to other people, even if they have the same symptoms that you have. It may harm them.

**Please read full Prescribing Information for [RYBREVANT \*FASPRO\*](#), [RYBREVANT](#), and [LAZCLUZE](#) available on this platform or at this presentation.**  
cp-493867v3

© Johnson & Johnson and its affiliates 2026 05/26 cp- 570727v2