

# Lexi's Story: Proactive Approaches

## ON-SCREEN TEXT

The following is a testimonial about a patient receiving RYBREVANT® (amivantamab-vmjw) at the time of filming. RYBREVANT® is indicated in combination with carboplatin and pemetrexed as a first-line treatment for people who have locally advanced or metastatic non—small cell lung cancer with EGFR exon 20 insertion mutations. Individual results may vary. Please talk with your doctor if you have any questions or to see if RYBREVANT® is right for you.

## Important Safety Information

RYBREVANT® is a prescription medicine used to treat adults with non—small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (*EGFR*) gene(s). RYBREVANT® can cause serious side effects such as infusion-related reactions and lung, blood clot, skin and eye problems. Please see additional Important Safety Information at the end of this video and full Prescribing Information available on this website.

## Lexi (VOICE OVER)

I was 37.

I was in great shape.

I was healthy.

I didn't didn't really think of lung cancer as something that could affect anyone.

## ON-SCREEN TEXT AND MUSIC

A RYBREVANT® Patient Story

Lexi

Lexi is a real patient and a volunteer with the SHARE Network, a Johnson & Johnson program made up of people who are dedicated to inspiring others through their personal health journeys and stories of caring. Every patient's experience is unique, and Lexi's journey is not representative of all patients. Lexi has been compensated for her time.

Lexi

We get a call from the oncologist, and he said, well, you know, it's not what we hoped. It's metastatic adenocarcinoma and, you know, consistent with primary lung cancer. And I was like, "What?" Like, I don't smoke. I've never smoked. And he was like, yeah, that doesn't matter. Hit me like a ton of bricks.

So, it was kind of like, you know, you just have a breath of a moment to swallow your grief and then go into action mode. And that's what I did. I ramped up on RYBREVANT® and chemo.

## ON-SCREEN TEXT AND MUSIC

Lexi's first experience with RYBREVANT®

The side effects really kind of came on slowly. Over time, I started developing a skin rash, paronychia, which is when you have an infection around your fingernails and it's really painful. It was just the new way of living.

## ON-SCREEN TEXT AND MUSIC

### Knowing when to listen to your body

I think that it's important to listen to your body. And so, breaks may need to happen and that's okay, and it can feel very stressful, and it can feel very unnerving.

So after I took a break in 2024, the summer of 2024, to kind of let my body recover, I had a PET scan September of 2024, and they found that the cancer regrew in my lung. At that time, my doctor wanted to try an oral drug trial. And so, I tried that for three months.

My doctor and I at that point decided that I had several options... we decided that surgery was a good next step for me. So I did a few rounds of chemo prior to surgery to prepare. And then I just was on chemo for a little while after surgery.

In May of 2025, we found growth in my brain and growth, regrowth in my iliac bone.

## ON-SCREEN TEXT AND MUSIC

### Reintroducing RYBREVANT® with a proactive approach

Because I had had such success with RYBREVANT prior, we decided that RYBREVANT was the best path forward, and there was in between when I had stopped taking it prior and the decision to take it again.

It was almost a year after I stopped taking it. There were new prophylactic measures that were recommended for treatment of side effects and new findings that came out around different things that someone could do to lessen the severity of the toxicities from the drug.

At that point, I was armed with so much more information and knowledge than I was before. I also didn't have a derm-, dermatology oncologist that I was working with the first time around proactively.

## ON-SCREEN TEXT AND MUSIC

### Tips to manage the treatment experience

There's certain type of treatments and therapies that are recommended that are found to significantly reduce the side effects. And it really has, in my case, significantly reduced the toxicity level to where I can live my life.

I take minocycline twice daily. It's a oral antibiotic and they either—doctors will either prescribe doxycycline or minocycline. It's a lot of little things that don't really take that much time out of your day to make a big difference you know, washing my hands or putting on creams to be able to live the rest of my day in a way that is more comfortable.

I think that it's a huge win to be able to take this drug, have access to this effective treatment, and have the ability to control symptoms to this degree is really remarkable.

## ON-SCREEN TEXT AND MUSIC

### Don't wait: treat proactively

You don't want to wait for these symptoms to come to try to combat them. I did that the first round, and treating them reactively is much more difficult to get them under control than if you treat them proactively.

Make sure that you and your doctor have these discussions, that you're on top of any type of new recommendations that come out and that you protect yourself.

People talk about shared decision-making, but it's really informed decision-making. There isn't a one-size-fits-all to what is right. There's no right or wrong. It's just about finding out what's important to you.

I keep seeing how fortunate I am to have the support system that I do, but I really am so fortunate. I have such good friends that I have, you know, from childhood, from college, post-college, family. Just I have the most amazing network of people, and they all want to do something.

It's okay to have moments where you're not strong and where it's acknowledging how hard it is. But you got to put that band-aid back on. You got to get up. You got to move. You got to find the things that make you happy and, and do those things and push through.

## ON-SCREEN TEXT AND MUSIC

Logo for RYBREVANT® (amivantamab-vmjw)  
Injection for IV Use | 350 mg/7 mL (50 mg/mL)

## ON-SCREEN TEXT AND NARRATOR

### **What are RYBREVANT *FASPRO* (amivantamab and hyaluronidase-lpuj) and RYBREVANT (amivantamab-vmjw)?**

RYBREVANT *FASPRO* and RYBREVANT are prescription medicines used to treat adults with non—small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s):

- in combination with LAZCLUZE (lazertinib), a prescription medicine, as a first-line treatment for NSCLC
- in combination with carboplatin and pemetrexed for NSCLC in people whose disease has worsened on or after treatment with an EGFR tyrosine kinase inhibitor (TKI)

- in combination with carboplatin and pemetrexed as a first-line treatment for NSCLC
- alone for the treatment of NSCLC in people whose disease has worsened on or after platinum-based chemotherapy.

Your healthcare provider will perform a test to make sure that RYBREVANT *FASPRO* or RYBREVANT is right for you.

It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE are safe and effective in children.

## **IMPORTANT SAFETY INFORMATION**

**Do not take RYBREVANT *FASPRO* if you are allergic to hyaluronidase or any of the ingredients in RYBREVANT *FASPRO*.**

**Before you receive RYBREVANT *FASPRO*, RYBREVANT or LAZCLUZE, tell your healthcare provider (HCP) about all of your medical conditions, including if you:**

- have a history of lung or breathing problems, other than cancer.
- are pregnant or plan to become pregnant. RYBREVANT *FASPRO*, RYBREVANT and LAZCLUZE can harm your unborn baby.

### **Females who are able to become pregnant:**

- Your HCP should do a pregnancy test before you start treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.
- If you are taking RYBREVANT *FASPRO* or RYBREVANT, use effective birth control (contraception) during treatment and for 3 months after your last dose. If you are taking LAZCLUZE, use effective birth control (contraception) during treatment and for 3 weeks after your last dose.
- Tell your HCP right away if you become pregnant or think you might be pregnant during treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

### **Males who have female partners who are able to become pregnant:**

- You should use effective birth control during treatment and for 3 weeks after your last dose of LAZCLUZE.
- are breastfeeding or plan to breastfeed. It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE pass into your breast milk. Do not breastfeed during treatment and for 3 months after your last dose of RYBREVANT *FASPRO* or RYBREVANT. Do not breastfeed during treatment and for 3 weeks after your last dose of LAZCLUZE.

**Tell your HCP about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. LAZCLUZE may affect the way other medicines work, and other medicines may affect how LAZCLUZE works. You should not start or stop any medicine before you talk with your HCP that prescribed LAZCLUZE.

### **What should I avoid while receiving RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE?**

RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause skin reactions. You should limit your time in the sun during and for 2 months after your treatment. Wear

protective clothing and use broad-spectrum sunscreen during treatment.

### **What are the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE?**

#### **RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE may cause serious side effects, including:**

- **allergic and injection-related reactions.** Allergic and injection-related reactions are common with RYBREVANT *FASPRO* and can be severe or serious. Tell your HCP right away if you get any of the following symptoms during or after your injection: shortness of breath, fever, chills, flushing, chest discomfort, dizziness or lightheadedness, or vomiting.
- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT but can be severe or serious and can include life-threatening (anaphylaxis) allergic reaction. Tell your HCP right away if you get any of the following symptoms during your infusion of RYBREVANT: shortness of breath, difficulty breathing, or wheezing, chest discomfort, swelling of your eyes, lips, or tongue, fever, chills, numbness of your tongue, lips, cheeks, or gums, flushing, skin rash, hives, or itching, nausea or vomiting, stomach cramps, lightheadedness, dizziness, or fainting, or headache.
- **lung problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause lung problems that can lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your HCP right away if you get any new or worsening lung symptoms, including shortness of breath, cough, or fever.
- **blood clot problems.** RYBREVANT *FASPRO* or RYBREVANT, when given in combination with LAZCLUZE, can cause blood clots in the veins of your legs (deep vein thrombosis) or lungs (pulmonary embolism) that can lead to death. Your HCP may start you on medicine to prevent blood clots for the first 4 months of treatment. Tell your HCP right away if you get any signs and symptoms of blood clots, including swelling, pain or tenderness in the leg, sudden unexplained chest pain, or shortness of breath.
- **skin problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause severe rash, including blisters, peeling, skin pain and sores, redness, raised acne-like bumps, itching, and dry skin. Your HCP may start you on an antibiotic for the first 3 months of treatment followed by an antibiotic lotion for your scalp for the next 9 months. During treatment, you should apply a non-comedogenic (does not clog pores) skin moisturizer (ceramide-based or other types of moisturizers that provide long lasting skin hydration and does not include drying ingredients) on your face and whole body (except scalp) and wash your hands and feet every day with 4% chlorhexidine solution. Tell your HCP right away if you get any skin reactions. Your HCP may treat you with a medicine(s) or send you to see a skin specialist (dermatologist) if you get skin reactions during treatment.
- **liver problems.** LAZCLUZE may cause severe liver problems, including increased liver function tests. Your HCP will do blood tests to check your liver before you start and during treatment. Tell your HCP right away if you get any symptoms of liver problems, including pain on the right side of your stomach area (abdomen), dark or brown (tea colored) urine, or yellowing of your skin or the white part of

your eyes.

- **eye problems.** RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause eye problems. Tell your HCP right away if you get symptoms of eye problems including eye pain, inflammation of eyelids, dry eyes, eye redness, blurred vision, changes in vision, itchy eyes, excessive tearing, and sensitivity to light. Your HCP may send you to see an eye specialist (ophthalmologist) if you get new or worsening eye problems during treatment. You should not use contact lenses until your eye symptoms are checked by an HCP.

**The most common side effects of RYBREVANT *FASPRO* in combination with LAZCLUZE include:** rash, infected skin around the nail, muscle and joint pain, feeling very tired, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, nausea, diarrhea, vomiting, constipation, decreased appetite, headache, and changes in certain blood tests.

**The most common side effects seen with intravenous (IV) RYBREVANT, which may be experienced with RYBREVANT *FASPRO*, are listed below:**

- **RYBREVANT in combination with LAZCLUZE:** rash, infected skin around the nail, infusion-related reaction, muscle and joint pain, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, unusual feeling in the skin (such as tingling or a crawling feeling), feeling very tired, diarrhea, constipation, COVID-19, bleeding, dry skin, decreased appetite, itchy skin, nausea, and changes in certain blood tests.
- **RYBREVANT in combination with carboplatin and pemetrexed:** rash, infected skin around the nail, infusion-related reaction, feeling very tired, nausea, sores in the mouth, constipation, swelling of hands, ankles, feet, face, or all of your body, decreased appetite, muscle and joint pain, vomiting, COVID-19, and changes in certain blood tests.
- **RYBREVANT given alone:** rash, infusion-related reaction, infected skin around the nail, muscle and joint pain, shortness of breath, nausea, swelling of hands, ankles, feet, face, or all of your body, cough, feeling very tired, sores in the mouth, constipation, vomiting, and changes in certain blood tests.

LAZCLUZE may cause fertility problems in males and females, which may affect your ability to have children. Talk to your HCP if this is a concern for you.

Your HCP may temporarily stop, decrease your dose, or completely stop your treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE if you have serious side effects.

These are not all of the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**General information about the safe and effective use of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE:**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LAZCLUZE for a condition for which it was not prescribed. Do not give LAZCLUZE to other people, even if they have the same

symptoms that you have. It may harm them.

**Please read full Prescribing Information for [RYBREVANT FASPRO](#), [RYBREVANT](#), and [LAZCLUZE](#) available on this platform or at this presentation.**

cp-493867v3

© Johnson & Johnson and its affiliates 2026 05/26 cp-570725v2