

Tania's Story: Moving Forward with Hope

ON-SCREEN TEXT

The following is a testimonial about a patient receiving RYBREVANT® (amivantamab-vmjw) at the time of filming. RYBREVANT® is indicated in combination with carboplatin and pemetrexed as a second-line treatment for people who have locally advanced or metastatic NSCLC with *EGFR* exon 19 deletions or exon 21 L858R substitution mutations, whose disease has progressed on or after treatment with an *EGFR* tyrosine kinase inhibitor. Individual results may vary. Please talk with your doctor if you have any questions or to see if RYBREVANT® is right for you.

Important Safety Information

RYBREVANT® is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (*EGFR*) gene(s). RYBREVANT® can cause serious side effects such as infusion-related reactions and lung, blood clot, skin and eye problems. Please see additional Important Safety Information at the end of this video and full Prescribing Information available on this website.

ON-SCREEN TEXT AND MUSIC

A RYBREVANT® Patient Story

Tania

Hi, I'm Tania, and I am from Northern California, just over the bridge from San Francisco in a county called Marin. And I am an entrepreneur and an inventor. I have one daughter, three brothers. They're all fantastic. We're very close.

Everybody in my family is a creative. Everybody has their own business. Yeah, I've always been like an outdoor person. I love to ski and I love to swim. I'm a big river girl in the summer. I did Taekwondo. I trained for years and years. I'm a second-degree black belt.

ON-SCREEN TEXT AND MUSIC

Tania is a real patient and a volunteer with the SHARE Network, a Johnson & Johnson program made up of people who are dedicated to inspiring others through their personal health journeys and stories of caring. Every patient's experience is unique and Tania's journey is not representative of all patients. Tania has been compensated for her time.

ON-SCREEN TEXT AND MUSIC

My Diagnosis

Tania

I was driving one morning, and all of a sudden, my sight got fractured, so I was seeing everything but in small, little multiples across my vision and I was like what the heck? It didn't last for more than like, I would say ten, fifteen seconds. It was super fast and it was just gone. And I was like, what was that?

I went to the ER and I saw this doctor and he was like, do you have any numbness anywhere, and I'm like, well, actually, I have, like, this little spot, like, on my jawline that feels like a little numb. And that alarmed him for some reason and was like, well, you know what, I'm going to send you in to an MRI and do a couple other little tests and we're going to see what it is.

And about an hour later, he came in and he said, yeah, you've got cancer, stage 4 lung cancer and you have 47 brain mets, 47 occurrences of cancer in your brain, and some on my liver and some on an adrenal, some on some, on a rib, some on the base of my back, like on a L2 or L3 on my spine.

They took a biopsy of the main tumor in my lung and sent it off for testing. I have *EGFR*, which is a type of lung cancer that happens to people that generally are non-smokers.

ON-SCREEN TEXT AND MUSIC

The Lung Cancer

Treatment Journey

Tania

They put me on an initial treatment, and that worked until it didn't. And in January of 2025 my doctor came back to me and said that she and her team would suggest that I start RYBREVANT along with a chemo. In the three-month and six-month markers since I started that treatment, the tumors that I had prior are gone. And so I have no more occurrence in my lung. It's gone from my adrenal and it's gone from the top of my liver.

ON-SCREEN TEXT AND MUSIC

This is Tania's personal experience.

Tania

My biggest side effect has been paronychia, which is where the cuticles around your nails become inflamed, and I have that on my toes. And I've been working with a podiatrist and with my oncologist on that, and bit by bit we've been managing that.

My podiatrist has me on a regimen between using a topical cleanser one week and then the next week I'll use a topical steroid.

I have a great team, a great oncology team. And for anybody that is facing cancer, I highly recommend that you put together a team and see what types of support you've got in your community. I have an oncology therapist, I have a palliative care person, I have a social worker and I also have a PT person that I can go to as well.

ON-SCREEN TEXT AND MUSIC

My Experience with Treatment

Tania

RYBREVANT and the chemo treatment that I'm on now is once every three weeks. I go into the infusion center at my local hospital and I am in a room for about six hours, six, six and a half hours.

ON-SCREEN TEXT AND MUSIC

RYBREVANT *FASPRO*[™]

This refers to how long the injection takes. Actual clinic time may vary. Your care team will also give you additional medications and take time to make sure you're doing well before you leave your visit.

Tania

Now that RYBREVANT *FASPRO* is available, it's going to be a game changer. It's going to change my time from hours to do the infusion to five-to-seven minutes for a injection. I can't wait to talk to my doctor about it.

ON-SCREEN TEXT AND MUSIC

This is Tania's personal experience.

Tania

When I got diagnosed, I knew that I could not go forward with running my business. I went through, like, a few months where I thought like huh, I don't have any creative thought. I'm usually thinking of, like, new ideas about stuff all the times, and I went through this period where I wasn't thinking of anything.

And then I just had this breakthrough where I went, well that's ridiculous. It's like, come up with something, do something. And it's going to be what it's going to be anyway. That was a big breakthrough for me where I realized that I had like a little bit of fear that was holding me down and I pushed through it.

I'm very excited about my creativity coming back. I've started training in Taekwondo again. I'm absolutely enjoying every single moment right now.

My outlook on life is super positive today. I'm very happy with my medical team and I'm grateful that there is so much work being done in the lung cancer research that a drug like RYBREVANT came out and that I'm able to be sitting here talking with you today. It's amazing. Because had this been 20 years ago, I might not be sitting here today.

The thing that really keeps me going day to day is just staying in the moment, staying present, and being appreciative for the life that I have right now.

ON-SCREEN TEXT AND MUSIC

RYBREVANT *FASPRO*[™] (amivantamab and hyaluronidase-lpuj)

RYBREVANT *FASPRO*[™] Prescribing Information, Horsham, PA: Janssen Biotech, Inc.
RYBREVANT[®] Prescribing Information, Horsham, PA: Janssen Biotech, Inc.

ON-SCREEN TEXT AND NARRATOR

What are RYBREVANT *FASPRO* (amivantamab and hyaluronidase-lpuj) and RYBREVANT (amivantamab-vmjw)?

RYBREVANT *FASPRO* and RYBREVANT are prescription medicines used to treat adults with non–small cell lung cancer (NSCLC) that has spread to other parts of the body (metastatic) or cannot be removed by surgery, and has certain abnormal epidermal growth factor receptor (EGFR) gene(s):

- in combination with LAZCLUZE (lazertinib), a prescription medicine, as a first-line treatment for NSCLC
- in combination with carboplatin and pemetrexed for NSCLC in people whose disease has worsened on or after treatment with an EGFR tyrosine kinase inhibitor (TKI)
- in combination with carboplatin and pemetrexed as a first-line treatment for NSCLC
- alone for the treatment of NSCLC in people whose disease has worsened on or after platinum-based chemotherapy.

Your healthcare provider will perform a test to make sure that RYBREVANT *FASPRO* or RYBREVANT is right for you.

It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE are safe and effective in children.

IMPORTANT SAFETY INFORMATION

Do not take RYBREVANT *FASPRO* if you are allergic to hyaluronidase or any of the ingredients in RYBREVANT *FASPRO*.

Before you receive RYBREVANT *FASPRO*, RYBREVANT or LAZCLUZE, tell your healthcare provider (HCP) about all of your medical conditions, including if you:

- have a history of lung or breathing problems, other than cancer.
- are pregnant or plan to become pregnant. RYBREVANT *FASPRO*, RYBREVANT and LAZCLUZE can harm your unborn baby.

Females who are able to become pregnant:

- Your HCP should do a pregnancy test before you start treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.
- If you are taking RYBREVANT *FASPRO* or RYBREVANT, use effective birth control (contraception) during treatment and for 3 months after your last dose. If you are taking LAZCLUZE, use effective birth control (contraception) during treatment and for 3 weeks after your last dose.
- Tell your HCP right away if you become pregnant or think you might be pregnant during treatment with RYBREVANT *FASPRO*, RYBREVANT, or

LAZCLUZE.

Males who have female partners who are able to become pregnant:

- You should use effective birth control during treatment and for 3 weeks after your last dose of LAZCLUZE.
- are breastfeeding or plan to breastfeed. It is not known if RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE pass into your breast milk. Do not breastfeed during treatment and for 3 months after your last dose of RYBREVANT *FASPRO* or RYBREVANT. Do not breastfeed during treatment and for 3 weeks after your last dose of LAZCLUZE.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. LAZCLUZE may affect the way other medicines work, and other medicines may affect how LAZCLUZE works. You should not start or stop any medicine before you talk with your HCP that prescribed LAZCLUZE.

What should I avoid while receiving RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE?

RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause skin reactions. You should limit your time in the sun during and for 2 months after your treatment. Wear protective clothing and use broad-spectrum sunscreen during treatment.

What are the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE?

RYBREVANT *FASPRO*, RYBREVANT, and/or LAZCLUZE may cause serious side effects, including:

- **allergic and injection-related reactions.** Allergic and injection-related reactions are common with RYBREVANT *FASPRO* and can be severe or serious. Tell your HCP right away if you get any of the following symptoms during or after your injection: shortness of breath, fever, chills, flushing, chest discomfort, dizziness or lightheadedness, or vomiting.
- **infusion-related reactions.** Infusion-related reactions are common with RYBREVANT but can be severe or serious and can include life-threatening (anaphylaxis) allergic reaction. Tell your HCP right away if you get any of the following symptoms during your infusion of RYBREVANT: shortness of breath, difficulty breathing, or wheezing, chest discomfort, swelling of your eyes, lips, or tongue, fever, chills, numbness of your tongue, lips, cheeks, or gums, flushing, skin rash, hives, or itching, nausea or vomiting, stomach cramps, lightheadedness, dizziness, or fainting, or headache.
- **lung problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause lung problems that can lead to death. Symptoms may be similar to those symptoms from lung cancer. Tell your HCP right away if you get any new or worsening lung symptoms, including shortness of breath, cough, or fever.
- **blood clot problems.** RYBREVANT *FASPRO* or RYBREVANT, when given in combination with LAZCLUZE, can cause blood clots in the veins of your legs (deep vein thrombosis) or lungs (pulmonary embolism) that can lead to death.

Your HCP may start you on medicine to prevent blood clots for the first 4 months of treatment. Tell your HCP right away if you get any signs and symptoms of blood clots, including swelling, pain or tenderness in the leg, sudden unexplained chest pain, or shortness of breath.

- **skin problems.** RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE can cause severe rash, including blisters, peeling, skin pain and sores, redness, raised acne-like bumps, itching, and dry skin. Your HCP may start you on an antibiotic for the first 3 months of treatment followed by an antibiotic lotion for your scalp for the next 9 months. During treatment, you should apply a non-comedogenic (does not clog pores) skin moisturizer (ceramide-based or other types of moisturizers that provide long lasting skin hydration and does not include drying ingredients) on your face and whole body (except scalp) and wash your hands and feet every day with 4% chlorhexidine solution. Tell your HCP right away if you get any skin reactions. Your HCP may treat you with a medicine(s) or send you to see a skin specialist (dermatologist) if you get skin reactions during treatment.
- **liver problems.** LAZCLUZE may cause severe liver problems, including increased liver function tests. Your HCP will do blood tests to check your liver before you start and during treatment. Tell your HCP right away if you get any symptoms of liver problems, including pain on the right side of your stomach area (abdomen), dark or brown (tea colored) urine, or yellowing of your skin or the white part of your eyes.
- **eye problems.** RYBREVANT *FASPRO*, RYBREVANT, and LAZCLUZE can cause eye problems. Tell your HCP right away if you get symptoms of eye problems including eye pain, inflammation of eyelids, dry eyes, eye redness, blurred vision, changes in vision, itchy eyes, excessive tearing, and sensitivity to light. Your HCP may send you to see an eye specialist (ophthalmologist) if you get new or worsening eye problems during treatment. You should not use contact lenses until your eye symptoms are checked by an HCP.

The most common side effects of RYBREVANT *FASPRO* in combination with LAZCLUZE include: rash, infected skin around the nail, muscle and joint pain, feeling very tired, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, nausea, diarrhea, vomiting, constipation, decreased appetite, headache, and changes in certain blood tests.

The most common side effects seen with intravenous (IV) RYBREVANT, which may be experienced with RYBREVANT *FASPRO*, are listed below:

- **RYBREVANT in combination with LAZCLUZE:** rash, infected skin around the nail, infusion-related reaction, muscle and joint pain, sores in the mouth, swelling of hands, ankles, feet, face, or all of your body, unusual feeling in the skin (such as tingling or a crawling feeling), feeling very tired, diarrhea, constipation, COVID-19, bleeding, dry skin, decreased appetite, itchy skin, nausea, and changes in certain blood tests.
- **RYBREVANT in combination with carboplatin and pemetrexed:** rash, infected skin around the nail, infusion-related reaction, feeling very tired, nausea, sores in the mouth, constipation, swelling of hands, ankles, feet, face, or all of your body, decreased appetite, muscle and joint pain, vomiting, COVID-19, and changes in

certain blood tests.

- **RYBREVANT given alone:** rash, infusion-related reaction, infected skin around the nail, muscle and joint pain, shortness of breath, nausea, swelling of hands, ankles, feet, face, or all of your body, cough, feeling very tired, sores in the mouth, constipation, vomiting, and changes in certain blood tests.

LAZCLUZE may cause fertility problems in males and females, which may affect your ability to have children. Talk to your HCP if this is a concern for you.

Your HCP may temporarily stop, decrease your dose, or completely stop your treatment with RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE if you have serious side effects.

These are not all of the possible side effects of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of RYBREVANT *FASPRO*, RYBREVANT, or LAZCLUZE:

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LAZCLUZE for a condition for which it was not prescribed. Do not give LAZCLUZE to other people, even if they have the same symptoms that you have. It may harm them.

Please read full Prescribing Information for [RYBREVANT *FASPRO*](#), [RYBREVANT](#), and [LAZCLUZE](#) available on this platform or at this presentation.

cp-493867v3

© Johnson & Johnson and its affiliates 2026 05/26 cp-570729v2